arXiv:1402.4919v4 [math.GR] 15 Mar 2014

Elementary p-adic Lie groups have finite constructible rank

Helge Glöckner

Abstract

The class of elementary totally disconnected groups is the smallest class of totally disconnected, locally compact, second countable groups which contains all discrete countable groups, all metrizable pro-finite groups, and is closed under extensions and countable ascending unions. To each elementary group G, a (possibly infinite) ordinal number $\mathrm{crk}(G)$ can be associated, its *constructible rank*. By a structure theorem of Phillip Wesolek, elementary p-padic Lie groups are among the basic building blocks for general σ-compact p-adic Lie groups. We characterize elementary p-adic Lie groups in terms of the subquotients needed to describe them. The characterization implies that every elementary p-adic Lie group has finite constructible rank.

Classification: primary 22E20; secondary 22E35, 22E46, 22E50

Key words: p-adic Lie group, series, topological simplicity, simple group, nearly simple group, Tits core, constructible group, scale function, rank, elementary group, centre, commutator group, subquotient, finiteness

Introduction and statement of the main results

By a theorem of Ph. Wesolek, every σ-compact p-adic Lie group G admits a series

$$\{1\} \lhd C \lhd S \lhd G$$

of closed subgroups such that G/S is finite, S/C is a finite direct product of non-discrete, compactly generated, topologically simple p-adic Lie groups, and C is elementary (cf. [12, Theorem 1.2]). In this sense, elementary p-adic Lie groups are among the basic building blocks for general σ-compact p-adic Lie groups. It is therefore desirable to improve the understanding of such groups.

We recall the definition of the constructible rank of an elementary group from [10]. Let Σ_0 be the class of all topological groups which are countable and discrete or metrizable and pro-finite. If $\lambda > 0$ is an ordinal number and Σ_μ has been defined for all $\mu < \lambda$, set $\Sigma_\lambda := \bigcup_{\mu<\lambda} \Sigma_\mu$ if λ is a limit ordinal. If λ has a precursor μ, let Σ_λ be the class of all topological groups

such that (a) G has a closed normal subgroup N such that N, $G/N \in \Sigma_\mu$, or (b) $G = \bigcup_{n\in\mathbb{N}} V_n$ for open subgroups $V_n \subseteq G$ such that $V_n \in \Sigma_\mu$ for all $n \in \mathbb{N}$. The constructible rank $\mathrm{crk}(G)$ of an elementary group G is defined as the minimum ordinal λ such that $G \in \Sigma_\lambda$.

Given a prime number p, let $\mathbb{Q}_p$ be the field of p-adic numbers and $\mathbb{Z}_p$ be the ring of p-adic integers. The p-adic Lie groups we consider are finite-dimensional analytic p-adic Lie groups as in [8]. We call a p-adic Lie group G *linear* if it admits an injective continuous homomorphism $G \to \mathrm{GL}_n(\mathbb{Q}_p)$ for some n. But linearity will not be assumed unless the contrary is stated. We say that a topological group G is *nearly simple* if every subnormal subgroup of G is open or discrete. If G is a p-adic Lie group, we write $L(G) := T_1(G)$ for its Lie algebra (the tangent space at the identity element $1 \in G$). As usual, $\mathrm{Ad}_G \colon G \to \mathrm{Aut}(L(G))$ is the adjoint representation, with kernel $\ker \mathrm{Ad}_G$.

We say that a p-adic Lie group G is *extraordinary* if G is nearly simple, $G = \ker \mathrm{Ad}_G$ (whence $L(G)$ is abelian) but the commutator group S' is non-discrete for each open subnormal subgroup $S \subseteq G$. Then G is non-discrete and non-abelian, in particular.

The author does not know whether extraordinary p-adic Lie groups exist (Remark 3.2). Linear Lie groups cannot be extraordinary (Remark 3.4).

Since discrete groups need not admit composition series, also p-adic Lie groups need not admit composition series (in terms of closed subnormal subgroups). Yet, they admit series which are close to such (Proposition 2.2). Our first main result is a structure theorem for general p-adic Lie groups, which describes the nearly simply subquotients needed to build up such groups.

Theorem A. *For every p-adic Lie group G, there exists a series*

$$G = G_0 \rhd G_1 \rhd \cdots \rhd G_n = \{1\}$$

of closed subgroups of G such that each of the subfactors $Q_j := G_{j-1}/G_j$ for $j \in \{1, \ldots, n\}$ has one of the following mutually exclusive properties:

(a) *Q_j is discrete*;

(b) *$Q_j \cong \mathbb{Z}_p$*;

(c) *Q_j is extraordinary*;

(d) *Q_j is isomorphic to a compact open subgroup of* $\mathrm{Aut}(\mathfrak{s})$ *for a simple p-adic Lie algebra* $\mathfrak{s}$; or

(e) *Q_j is topologically simple, compactly generated, non-discrete, and isomorphic to an open subgroup of* $\mathrm{Aut}(\mathfrak{s})$ *for a simple p-adic Lie algebra* $\mathfrak{s}$.

We also obtain the following characterization:

Theorem B. *A p-adic Lie group G is elementary if and only if it admits a series*

$$G = G_0 \rhd G_1 \rhd \cdots \rhd G_n = \{1\}$$

of closed subgroups of G such that each of the subfactors $Q_j := G_{j-1}/G_j$ for $j \in \{1, \ldots, n\}$ has one of the following mutually exclusive properties:

(a) *Q_j is countable and discrete*;

(b) *$Q_j \cong \mathbb{Z}_p$*;

(c) *Q_j is extraordinary and σ-compact*; *or*

(d) *Q_j is isomorphic to a compact open subgroup of* $\mathrm{Aut}(\mathfrak{s})$ *for a simple p-adic Lie algebra* $\mathfrak{s}$.

The preceding characterization implies:

Theorem C. *Every elementary p-adic Lie group has finite constructible rank.*

In fact, if G is such a group and $G = G_0 \rhd G_1 \rhd \cdots \rhd G_n = \{1\}$ a series as in Theorem B, then $\mathrm{crk}(G) \leq n + 2e$, where $e \in \mathbb{N}_0$ is the number of indices $j \in \{1, \ldots, n\}$ for which G_{j-1}/G_j is extraordinary.

We also provide a number of illustrating examples and some related results for Lie groups over other fields.

1 Terminology, notation and basic facts

We shall write $\mathbb{N} := \{1, 2, \ldots\}$ and $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$. If G is a group, we write G' for its commutator group and $G^{(n)} := (G^{(n-1)})'$ for the terms of the derived series, with $G^{(0)} := G$. We write $\mathrm{Aut}(\mathfrak{g})$ for the group of automorphisms of a Lie algebra $\mathfrak{g}$, and define $\mathrm{ad}(x)\colon \mathfrak{g} \to \mathfrak{g}$, $y \mapsto [x, y]$ for $x \in \mathfrak{g}$. Recall the

definition of the adjoint representation: If G is a p-adic Lie group and $g \in G$, one considers the inner automorphism $I_g \colon G \to G$, $I_g(x) := gxg^{-1}$ and defines $\mathrm{Ad}_G(g) := L(I_g) := T_1(I_g)$ as the associated Lie algebra automorphism (the tangent map of I_g at 1). Then $\mathrm{Ad}_G \colon G \to \mathrm{Aut}(L(G))$ is continuous and $L(\mathrm{Ad}_G)(x) = \mathrm{ad}(x)$ for each $x \in L(G)$(see [3]). Thus $\mathrm{ad}(x) = 0$ for all $x \in \mathfrak{g}$ and thus $L(G)$ is abelian whenever $\mathrm{Ad}_G(g) = \mathrm{id}_{\mathfrak{g}}$ for all $g \in G$ (e.g., if G is extraordinary). As usual, a topological group G is called *compactly generated* if it has a compact generating set. It is called *topologically simple* if $G \neq \{1\}$ and G does not have closed normal subgroups other than $\{1\}$ and G. An automorphism of a topological group G is a group automorphism which is a homeomorphism. A subgroup $H \subseteq G$ is called *topologically characteristic* if $\alpha(H) = H$ for each automorphism of the topological group G. The *radical* $R(G)$ of a (Hausdorff) topological group G is defined as the union of all closed, normal, soluble subgroups.

We shall use a fact which is immediate from classical work by H. Bass:

Lemma 1.1 *Let $\mathbb{K}$ be a totally disconnected local field, $\mathbb{O} \subseteq \mathbb{K}$ be the maximal compact subring, $n \in \mathbb{N}$ and $G \subseteq \mathrm{GL}_n(\mathbb{K}))$ be a subgroup with the following properties:*

(a) *Every $g \in G$ generates a relatively compact subgroup of $\mathrm{GL}_n(\mathbb{K})$;*

(b) *If $V \subseteq \mathbb{K}^n$ is a G-invariant vector subspace, then $V = \mathbb{K}^n$ or $V = \{0\}$.*

Then there exists $x \in \mathrm{GL}_n(\mathbb{K})$ such that $xGx^{-1} \subseteq \mathrm{GL}_n(\mathbb{O})$.

Proof. Let $\mathrm{tr} \colon M_n(\mathbb{K}) \to \mathbb{K}$ be the trace on the algebra of $n \times n$-matrices, and $|.|$ be an ultrametric absolute value on $\mathbb{K}$ defining its topology. For each $g \in G$, the subgroup $K := \overline{\langle g \rangle}$ of $\mathrm{GL}_n(\mathbb{K})$ is compact by hypothesis. Hence K is conjugate to a subgroup of $\mathrm{GL}_n(\mathbb{O})$ (see Theorem 1 in Appendix 1 to [8, Part II, Chapter IV]), say $xKx^{-1} \subseteq \mathrm{GL}_n(\mathbb{O})$. As a consequence, $|\mathrm{tr}(g) = |\mathrm{tr}(xgx^{-1})| \leq 1$. Thus $\mathrm{tr}(G)$ is bounded, whence G is relatively compact by [1, Corollary 1.3 (c)]. Therefore G is conjugate to a subgroup of $\mathrm{GL}_n(\mathbb{O})$ (using [8] again). □

Lemma 1.2 *For every p-adic Lie group G, its radical $R(G)$ is a closed, topologically characteristic subgroup of G whose Lie algebra $L(R(G))$ is contained in the radical $\mathfrak{r}$ of $L(G)$. Moreover, G has a soluble closed normal subgroup N which is open in $R(G)$.*

Proof. If N_1 and N_2 are soluble, closed normal subgroups of G, then $\overline{N_1N_2}$ is soluble as $N_1N_2/N_2 \cong N_1/(N_1 \cap N_2)$ and N_2 are soluble. Hence also $\overline{N_1N_2}$ is soluble (see [3, Chapter 3, §9, no. 1, Corollary 1]). The set $\mathcal{S}$ of soluble, closed normal subgroups of G is therefore directed under inclusion. Since every p-adic Lie group has a compact, open subgroup U which satisfies an ascending chain condition on closed subgroups, $N \cap U$ must become stationary, showing that $U \cap \bigcup_{N \in \mathcal{S}} N$ is closed. Being locally closed by the preceding, the subgroup $R(G) := \bigcup_{N \in \mathcal{S}} N$ of G is closed. It is immediate from the definition that $R(G)$ is topologically characteristic. If we choose $N \in \mathcal{S}$ such that $U \cap R(G) = U \cap N$, then N is open in $R(G)$. Hence $L(R(G)) = L(N)$, which is a soluble ideal (and hence contained in $\mathfrak{r}$) as N is normal and soluble (cf. [3, Chapter III, §9, no. 2, Proposition 3]). □

We also need a variant of [5, Lemma 6.4].[1]

Lemma 1.3 *Let G be a linear p-adic Lie group. Then the radical $R(G)$ is soluble and its Lie algebra $L(R(G))$ coincides with the radical $\mathfrak{r}$ of the Lie algebra $L(G)$.*

Proof. We may assume that $G \subseteq \mathrm{GL}_n(\mathbb{Q}_p)$ for some n (such that the Lie group topology on G is finer than the induced topology). Using the next lemma, we can copy the proof of [5, Lemma 6.4] to see that $L(R(G)) = \mathfrak{r}$. For every soluble closed normal subgroup $N \subseteq G$, the closure $\overline{N}$ in $\mathrm{GL}_n(\mathbb{Q}_p)$ is normal in $\overline{G}$ and soluble. Hence $R(G) \subseteq R(\overline{G})$. Since $R(\overline{G})$ is soluble (see [5, p. 20]), we deduce that also $R(G)$ is soluble. □

Lemma 1.4 *Let $\mathbb{K}$ be an infinite field, $n \in \mathbb{N}$ and $G \subseteq \mathrm{GL}_n(\mathbb{K})$ be a subgroup, endowed with a topology making it a topological group [2] which has an open subgroup which is soluble (resp., nilpotent). Then G has an open* normal *subgroup which is soluble (resp., nilpotent).*

Proof. We can copy the proof of [5, Lemma 6.1]. □

Remark 1.5 As we do not restrict our attention to p-adic Lie groups which are linear, we have to face certain difficulties:

[1] The lemma in [5] only tackles p-adic Lie groups that are closed subgroups of $\mathrm{GL}_n(\mathbb{Q}_p)$.

[2] The proof only uses that all inner automorphisms of G are continuous.

(a) The radical of a p-adic Lie group need not be soluble. For example, for each $n \in \mathbb{N}$ we can choose a soluble group H_n with $(H_n)^{(n)} \neq \{1\}$ and define G as the group $\bigoplus_{n\in\mathbb{N}} H_n$ of all $(h_n)_{n\in\mathbb{N}} \in \prod_{n\in\mathbb{N}} H_n$ such that $h_n = 1$ for all but finitely many n (endowed with the discrete topology).

(b) If G has a closed normal subgroup N such that $R(N)$ is not soluble, then it is not clear whether $R(N) \subseteq R(G)$.

(c) If $R(G)$ is not soluble, then it is not clear whether $G/R(G)$ has trivial radical.

(d) If G is not linear, then we do not know whether $L(R(G))$ must coincide with the radical of $L(G)$. If extraordinary p-adic Lie groups exist, then the answer is negative.

1.6 The class of elementary totally disconnected groups has the following permanence properties: *Closed subgroups of elementary groups and Hausdorff quotients of elementary groups are elementary* [11, Theorems 1.2 and 1.3].

2 Structural results on p-adic Lie groups

In this section, we provide auxiliary results on p-adic Lie groups which will help us to prove the main theorems.

Definition 2.1 We say that a topological group G is *nearly simple* if every closed subnormal subgroup $S \subseteq G$ is open in G or discrete.

In this section, we prove four propositions. The first one provides series which can later be refined further to obtain Theorems A and B.

Proposition 2.2 *Every p-adic Lie group G admits a series*

$$G = G_0 \rhd G_1 \rhd \cdots \rhd G_n = \{1\}$$

of closed subgroups G_j such that G_{j-1}/G_j is nearly simple for all $j \in \{1, \ldots, n\}$.

The second proposition provides information on nearly simple groups.

Proposition 2.3 *Let G be a p-adic Lie group, with Lie algebra $\mathfrak{g} := L(G)$. If G is nearly simple, then exactly one of the following cases occurs:*

(a) *G is discrete;*

(b) $\dim(G) = 1$ *and G has an open subnormal subgroup W such that $W/D \cong \mathbb{Z}_p$ for some discrete normal subgroup $D \subseteq W$;*

(c) *G has an open normal subgroup which is extraordinary;*

(d) *The Lie algebra $\mathfrak{g}$ is simple and $\mathrm{Ad}_G \colon G \to \mathrm{Aut}(\mathfrak{g})$ has discrete kernel. In this case, $Q := G/\ker \mathrm{Ad}_G$ is a nearly simple group such that each closed, non-trivial subnormal subgroup of Q is open in Q and $\mathrm{Ad}_Q \colon Q \to \mathrm{Aut}(\mathfrak{g})$ is injective.*

Further information on the groups Q from part (d) of Proposition 2.3 is provided by the next proposition.

Proposition 2.4 *Let G be a p-adic Lie group with simple Lie algebra $\mathfrak{g} := L(G)$. If G is nearly simple and $\mathrm{Ad}_G \colon G \to \mathrm{Aut}(\mathfrak{g})$ is injective, then exactly one of the cases* (a) *and* (b) *occurs:*

(a) *G is isomorphic to a compact open subgroup in $\mathrm{Aut}(\mathfrak{g})$;*

(b) *G has an open, topologically characteristic, non-compact subgroup M which is topologically simple, non-discrete, and compactly generated.*

If, moreover, G is elementary, then (a) *holds.*

The next proposition provides vital information on extraordinary groups.

Proposition 2.5 *Let G be an extraordinary p-adic Lie group. Then $G = \bigcup_{n\in\mathbb{N}} G_n$ with open subgroups $G_1 \subseteq G_2 \subseteq \cdots$ such that each G_n has an open centre. Hence, if G is σ-compact, then G is elementary with $\mathrm{crk}(G) \leq 2$.*

Proof. Since $\mathrm{Ad}_G(g) = \mathrm{id}_{L(G)}$, each $g \in G$ centralizes an open subgroup of G. Let $V_1 \supseteq V_2 \supseteq \cdots$ be compact open subgroups which form a base of identity neighbourhoods in G. Let $C_G(V_n)$ be the centralizer of V_n in G. Then $C_G(V_1) \subseteq C_G(V_2) \subseteq \cdots$ and

$$G = \bigcup_{n\in\mathbb{N}} C_G(V_n) \tag{1}$$

by the preceding. By Baire's category theorem, $C_G(V_N)$ is open in G for some N. After passage to a subsequence, we may therefore assume that

$C_G(V_n)$ is open in G for each $n \in \mathbb{N}$. Note that $W_n := V_n \cap C_G(V_n)$ is a compact open subgroup of $C_G(V_n)$ which is contained in the centre $Z(C_G(V_n))$. Hence $C_G(V_n)$ has open centre. If G is σ-compact, then $C_G(V_n)$ is elementary with constructible rank ≤ 1, since the compact open subgroup W_n is central (and hence normal) in $C_G(V_n)$. Hence $\mathrm{crk}(G) \leq 2$, by (1). □

Several lemmas and observations will help us to prove Propositions 2.2–2.4.

Lemma 2.6 *If a topological group G is nearly simple, then also each closed subnormal subgroup $S \subseteq G$ is nearly simple.*

Proof. If T is a closed subnormal subgroup of S, then T also is a closed subnormal subgroup of G and hence open in G (in which case it is also open in S) or discrete. □

Lemma 2.7 *If V is an open subgroup of a topological group G and V is nearly simple, then also G is nearly simple.*

Proof. If $S \subseteq G$ is a closed subnormal subgroup of G, then $S \cap V$ is a closed subnormal subgroup of V. Since V is nearly simple, $S \cap V$ is open in V or discrete. If $S \cap V$ is open in V, then S is open in G. If $S \cap V$ is discrete, then S is discrete. □

Lemma 2.8 *Let G be a topological group and $D \subseteq G$ be a discrete normal subgroup. If G is nearly simple, then also G/D is nearly simple.*

Proof. Let $q\colon G \to G/D$ be the canonical quotient map. If S is a closed subnormal subgroup of G/D, then $q^{-1}(S)$ is a closed subnormal subgroup of G and hence open (in which case also $S = q(q^{-1}(S))$ is open) or discrete, in which case also $S \cong q^{-1}(S)/D$ is discrete. □

Lemma 2.9 *Let G be a topological group and $S \subseteq G$ be a closed subnormal subgroup. Then there exists a series $G = G_0 \rhd G_1 \rhd \cdots \rhd G_k = S$ such that G_j is closed in G for all $j \in \{1, \ldots, k\}$.*

Proof. If $S \subseteq G$ is a closed subnormal subgroup, we find a series $G = G_0 \rhd G_1 \rhd \cdots \rhd G_k = S$. After replacing each G_j with its closure, we may assume without loss of generality that all of $G_0, \ldots, G_k$ are closed. □

Lemma 2.10 *If a p-adic Lie group G has dimension $\dim(G) \leq 1$ or its Lie algebra $\mathfrak{g} := L(G)$ is simple, then G is nearly simple.*

Proof. If $\dim(G) = 0$, then G (and hence every subgroup of G) is discrete.

If $\dim(G) = 1$, then every closed subgroup of G has dimension 1 (in which case it is open) or dimension 0 (in which case it is discrete).

If $\mathfrak{g}$ is a simple Lie algebra and $N \subseteq G$ a non-discrete closed normal subgroup, then $L(N)$ is a non-zero ideal in $\mathfrak{g}$ and thus $L(N) = \mathfrak{g}$, entailing that N is open in G. If $S \subseteq G$ is a closed subnormal subgroup, we find a series $G = G_0 \rhd G_1 \rhd \cdots \rhd G_k = S$ with closed subgroups $G_0, \ldots, G_k$, by Lemma 2.9. Then all of $G_0, \ldots, G_k$ are nearly simple, by Lemma 2.6. By the preceding, G_1 is open in G and thus $L(G_1) = \mathfrak{g}$ is simple. Repeating the argument, we see that G_j is open in G_{j-1} for all $j \in \{1, \ldots, k\}$. In particular, $S = G_k$ is open in G. □

Proof of Proposition 2.2. If G is nearly simple (e.g., if $\dim(G) \leq 1$), we can take $G_0 := G$ and $G_1 := \{1\}$.

Induction step. Assume that $\dim(G) \geq 2$ and G is not nearly simple. Then there exists a closed subnormal subgroup S of G which is neither open in G nor discrete. Thus $0 < \dim(S) < \dim(G)$. Since S is subnormal, we find a series $G = G_0 \rhd G_1 \rhd \cdots \rhd G_k = S$ of closed subgroups (see Lemma 2.9). We may assume that S has been chosen such that $\dim(S)$ is maximal. Hence $G_1, \ldots, G_{k-1}$ are open in G and thus G_{j-1}/G_j is discrete (and hence nearly simple) for all $j \in \{1, \ldots, k-1\}$. Also $Q := G_{k-1}/G_k$ is nearly simple. To see this, let $q\colon G_{k-1} \to Q$ be the canonical quotient map. If the assertion was false, we could find a closed subnormal subgroup $T \subseteq Q$ such that $0 < \dim(T) < \dim(Q)$. Thus $R := q^{-1}(T)$ would be a closed subnormal subgroup of G with $\dim(G_k) < \dim(T) < \dim(G)$, contradicting the maximality of $\dim(S)$. Since $\dim(G_k) < \dim(G)$, by induction we have a series $G_k \rhd G_{k+1} \rhd \cdots \rhd G_n = \{1\}$ of closed subgroups with nearly simple quotients. □

Proof of Proposition 2.3. If G is discrete, we have (a). Now assume that $\dim(G) \geq 1$.

Case 1: *G has an open subnormal subgroup S with discrete commutator group S'.* Then $Q := S/S'$ is an abelian Lie group with $L(Q) = \mathfrak{g}$. By Lemmas 2.6 and 2.8, Q is nearly simple. Let $\mathfrak{a} \subseteq \mathfrak{g}$ be a 1-dimensional vector subspace and $\gamma\colon \mathbb{Z}_p \to Q$ be a continuous homomorphism with $0 \neq \gamma'(0) \in \mathfrak{a}$. Then $A := \gamma(\mathbb{Z}_p)$ is a compact Lie subgroup of Q with $L(A) = \mathfrak{a}$ and $\gamma\colon \mathbb{Z}_p \to$

A is an isomorphism. Since A is normal and non-discrete, A is open in Q and so $\mathfrak{g} = \mathfrak{a}$ is 1-dimensional. Let $q\colon S \to Q$ be the quotient map. Then $W := q^{-1}(A)$ is open and subnormal in G and $W/S' \cong \mathbb{Z}_p$, i.e., (b) holds.

Case 2: *Each open subnormal subgroup of G has non-discrete commutator group.* We then consider $\mathrm{Ad}_G\colon G \to \mathrm{Aut}(G)$ and note that $\ker \mathrm{Ad}_G$ is a closed normal subgroup of G. Hence either $\ker \mathrm{Ad}_G$ is open or $\ker \mathrm{Ad}_G$ is discrete. If $S := \ker \mathrm{Ad}_G$ is open, then $\mathrm{Ad}_S(g) = \mathrm{Ad}_G(g) = \mathrm{id}_{\mathfrak{g}}$ for all $g \in S$ and thus S is a nearly simple, extraordinary, open normal subgroup of G (whence we have (c)). Now assume that $\ker \mathrm{Ad}_G$ is discrete. Then $Q := G/\ker \mathrm{Ad}_G$ is nearly simple by Lemma 2.8. Moreover, $R(Q)$ is discrete. To see this, suppose that the closed normal subgroup $R(Q)$ of Q was non-discrete (and hence open). Then Q has a soluble, open, normal subgroup N (Lemma 1.2). Choose $k \in \mathbb{N}_0$ maximal such that $N^{(k)}$ is non-discrete. Let $q\colon G \to Q$ be the quotient map. Then $H := q^{-1}(\overline{N^{(k)}})$ is a non-discrete, closed normal subgroup of G (and thus open) such that H' is discrete (which contradicts the hypothesis of Case 2). Thus indeed $R(Q)$ is discrete. Since $L(Q) = \mathfrak{g}$ and $\mathrm{Ad}_Q \circ q = \mathrm{Ad}_G$, where q is surjective and the kernel is $\ker \mathrm{Ad}_G$ only, we deduce that Ad_Q is injective. In particular, Q is a *linear* p-adic Lie group. Let $\mathfrak{r}$ be the radical of $\mathfrak{g} = L(Q)$. Since Q is a linear p-adic Lie group, the discreteness of $R(G)$ implies $\mathfrak{r} = \{0\}$ (see Lemma 1.3). Thus $\mathfrak{g}$ is semisimple. If $S \subseteq Q$ is a non-trivial closed subnormal subgroup, then Lemma 2.9 provides a series

$$Q = Q_0 \rhd Q_1 \rhd \cdots \rhd Q_k = S$$

of closed subgroups of Q. Then Q_j is open in Q or discrete, for each $j \in \{0, \ldots, k\}$. We show by induction on j that Q_j is open in Q. Since $Q_0 = Q$, this is trivial if $j = 0$. If $j > 0$ and Q_{j-1} is open in Q, let us show that also Q_j is open in Q. If not, then Q_j was discrete and thus $P := Q_{j-1}/Q_j$ would be a group with Lie algebra $\mathfrak{g}$. Let $p\colon Q_{j-1} \to P$ be the quotient map. Then $\mathrm{Ad}_P \circ p = \mathrm{Ad}_{Q_{j-1}}$ where $\mathrm{Ad}_{Q_{j-1}} = \mathrm{Ad}_Q|_{Q_{j-1}}$ is injective. As a consequence, p must be injective and thus $Q_j = \{1\}$, contrary to $Q_j \supseteq S \neq \{1\}$. Hence Q_j is open in Q and thus also $S = Q_k$ is open, by induction.

It only remains to show that $\mathfrak{g}$ is simple. Write $\mathfrak{g} = \mathfrak{s}_1 \oplus \cdots \oplus \mathfrak{s}_m$ with simple ideals $\mathfrak{s}_1, \ldots, \mathfrak{s}_m$. Let $W \subseteq \mathrm{Aut}(\mathfrak{g})$ be the group of all $\phi \in \mathrm{Aut}(\mathfrak{g})$ such that $\phi(\mathfrak{s}_j) = \mathfrak{s}_j$ for all $j \in \{1, \ldots, m\}$. Then W is a normal subgroup of finite index in $\mathrm{Aut}(\mathfrak{g})$ (complemented by a finite group of permutations). As a consequence, $V := \mathrm{Ad}_Q^{-1}(W)$ is an open normal subgroup in Q. We identify W with $\mathrm{Aut}(\mathfrak{s}_1) \times \cdots \times \mathrm{Aut}(\mathfrak{s}_m)$ and let $\pi_j\colon W \to \mathrm{Aut}(\mathfrak{s}_j)$ be the

canonical projection. Let exp be an exponential function for V (see [3]). If we had $m \geq 2$, we would have

$$\mathrm{Ad}_V(\exp(x))(y) = e^{\mathrm{ad}(x)}(y) \in y + [\mathfrak{s}_j, \mathfrak{s}_k] = \{y\}$$

for all x in a 0-neighbourhood in $\mathfrak{s}_j$ and all $y \in \mathfrak{s}_k$, showing that

$$(\pi_k \circ \mathrm{Ad}_V)(\exp(x)) = \mathrm{id}_{\mathfrak{s}_k}$$

for all x in some identity neighbourhood in $\mathfrak{s}_j$. Thus $\ker(\pi_k \circ \mathrm{Ad}_V)$ is open in V for all $k \in \{1, \ldots, m\}$. But then $\bigcap_{k=1}^m \ker(\pi_k \circ \mathrm{Ad}_V) = \ker \mathrm{Ad}_V$ is open, contradiction. Hence $m = 1$ and thus $\mathfrak{g} = \mathfrak{s}_1$ is simple. We have all of (d). □

Proof of Proposition 2.4. Since $\mathrm{Ad}_G \colon G \to \mathrm{Aut}(\mathfrak{g})$ is an injective continuous homomorphism and $L(\mathrm{Aut}(\mathfrak{g})) \cong \mathfrak{g}$, we see that Ad_G is étale (i.e., a local diffeomorphism). Being also injective, Ad_G is an isomorphism of Lie groups onto an open (and hence closed) subgroup of $\mathrm{Aut}(\mathfrak{g})$. Let $G^\dagger$ be the Tits core of G, i.e., the subgroup generated by the contraction groups

$$U_g := \{x \in G \colon \ g^n x g^{-n} \to 1 \text{ as } n \to +\infty \ \}$$

for $g \in G$ (which are closed since G is a p-adic Lie group [9]), as in [4]. Then $G^\dagger$ is topologically characteristic in G and hence so is its closure $M := \overline{G^\dagger}$. In particular, M is normal in G. Hence M is open in G or $M = \{1\}$ (see Proposition 2.3 (d)).

Case 1: $M = \{1\}$. Then $G^\dagger = \{1\}$, whence $U_g = \{1\}$ for each $g \in G$ and thus $s_G(g) = 1$, i.e., the scale function $s_G \colon G \to \mathbb{N}$ (as defined in [13]) is identically 1 (cf. [2, Proposition 3.21 (3)]). Then each eigenvalue of $\mathrm{Ad}_G(g)$ in an algebraic closure of $\mathbb{Q}_p$ has absolute value 1 (see [6, Corollary 3.6]). This entails that $\mathrm{Ad}_G(g)$ is an isometry for some choice of ultrametric norm on $\mathfrak{g}$ (see Lemma 3.3 in [6] and its proof). As a consequence, the subgroup generated by $\mathrm{Ad}_G(g)$ is relatively compact. Since $\mathrm{Ad}_G(G)$ is closed, we deduce with Lemma 1.1 that $\mathrm{Ad}_G(G)$ is compact. We have (a).

Case 2: Assume that M is open in G. Then $M \neq \{1\}$. Since each non-trivial contraction group U_g is non-discrete, also M is non-discrete. We show that M is topologically simple. If $N \subseteq M$ is a non-trivial closed normal subgroup, then N is subnormal in G. Hence N is open in G. If $g \in G$ and $x \in U_g$, then $g^n x g^{-n} \in N$ for some $n \in \mathbb{N}$. Thus $x \in g^{-n} N g^n = N$ (using that N is normal). Thus $G^\dagger \subseteq N$ and thus $M \subseteq N$ (as N is closed). Thus $N = M$

and topological simplicity is established. Finally, by [5, Proposition 6.5], there exists a group $R := S(\mathbb{Q}_p)$ of $\mathbb{Q}_p$-rational points of a simple isotropic $\mathbb{Q}_p$-algebraic group S such that $M \cong (R^\dagger)^n/Z$ for some $n \in \mathbb{N}$ and closed central subgroup $Z \subseteq (R^\dagger)^n$. Since $R^\dagger$ is compactly generated [7, Chapter I, Corollary 2.3.5], also M is compactly generated.

To complete the proof, recall that an elementary group that is topologically simple and compactly generated must be discrete [10, Theorem 4.6]. If M is as in (b) and G is elementary, then also the closed subgroup M is elementary (see 1.6). Since M is non-discrete, this contradicts the preceding fact. □

3 Proofs for Theorems A–C

Proof of Theorem A and B. We may assume that G is non-discrete. By Proposition 2.2, G admits a series $G = H_0 \rhd H_1 \rhd \cdots \rhd H_m = \{1\}$ of closed subgroups such that H_{j-1}/H_j is nearly simple for all $j \in \{1, \ldots, m\}$. The proof is by induction on $m \in \mathbb{N}$. If $m = 1$, then G is nearly simple. Case 1: If $L(G)$ is abelian, then Proposition 2.3 (b) or (c) yields a series of the desired form. Case 2: If $L(G)$ is simple, set $Q := G/\ker \mathrm{Ad}_G$. Then Propositions 2.3 (d) and 2.4 show that $\ker \mathrm{Ad}_G$ is discrete and either

(i): Q is isomorphic to a compact open subgroup in $\mathrm{Aut}(\mathfrak{g})$ (in which case $G \rhd \ker \mathrm{Ad}_G \rhd \{1\}$ is as desired); or

(ii): Q has an open characteristic subgroup M as in Proposition 2.4 (b), in which case we let $q\colon G \to Q$ be the canonical quotient map and take $G \rhd q^{-1}(M) \rhd \ker \mathrm{Ad}_G \rhd \{1\}$.

By Prop. 2.4, (ii) cannot occur if G is elementary, as Q is elementary by 1.6.

Induction step. If $m > 1$, we have a series $G/H_1 = P_0 \rhd P_1 \rhd \cdots \rhd P_k = \{1\}$ of the desired form (by the case $m = 1$). We let $q\colon G \to G/H_1$ be the quotient map and define $G_j := q^{-1}(P_j)$ for $j \in \{0, \ldots, k\}$. Also, by the case $m - 1$ we have a series $H_1 = G_k \rhd G_{k+1} \cdots \rhd G_n = \{1\}$ of the desired form, for some $n > k$. Then $G = G_0 \rhd G_1 \rhd \cdots \rhd G_0 = \{1\}$ is as desired. □

Lemma 3.1 *Let G be an elementary group and $G = G_0 \rhd G_1 \rhd \cdots \rhd G_n = \{1\}$ be a series of closed subgroups. Then G_j and $Q_j := G_{j-1}/G_j$ are elementary*

for all $j \in \{1, \ldots, n\}$,

$$\operatorname{crk}(G_{j-1}) \leq \max\{\operatorname{crk}(Q_j), \operatorname{crk}(G_j)\} + 1 \quad \textit{for all } j = n-1, n-2, \ldots, 1, \tag{2}$$

and $\operatorname{crk}(G_{n-1}) = \operatorname{crk}(Q_n)$. *In particular,* $\operatorname{crk}(G)$ *is finite if* $\operatorname{crk}(Q_j)$ *is finite for all* $j \in \{1, \ldots, n\}$.

Proof. G_j and Q_j are elementary by 1.6. The remaining assertions are clear from the definition of the constructible rank. □

Proof of Theorem C. Choose a series $G = G_0 \rhd G_1 \rhd \cdots \rhd G_n = \{1\}$ as in Theorem B, and set $Q_j := G_{j-1}/G_j$ for $j \in \{1, \ldots, n\}$. Then $\operatorname{crk}(Q_j) = 0$ if Q_j is discrete or compact (which holds in the cases (a), (b) and (d) of Theorem B). Otherwise (in case (c) of Theorem B), Q_j is extraordinary and thus (since Q_j is elementary), $\operatorname{crk}(Q_j) \leq 2$ by Proposition 2.5. Let e be the number of indices j such that G_j is extraordinary. Combining the preceding estimates for $\operatorname{crk}(Q_j)$ with Lemma 3.1, we obtain $\operatorname{crk}(G) \leq n + 2e$, In particular, $\operatorname{crk}(G)$ is finite. □

Remark 3.2 The author does not know whether extraordinary p-adic Lie groups exist. For which dimensions do they exist? For which dimensions can we have extraordinary p-adic Lie groups which are, moreover, σ-compact and hence elementary?

Remark 3.3 If a p-adic Lie group G is extraordinary in the sense defined above, then the radical $R(S)$ is discrete for each open subnormal subgroup S of G (and thus G is extraordinary in a weaker sense, as used in version v2 of this preprint). In fact, if $R(S)$ was non-discrete (and thus open), we could choose a soluble, open, normal subgroup N of S (by Lemma 1.2). Choose $k \in \mathbb{N}_0$ maximal such that $N^{(k)}$ is non-discrete. Then $H := \overline{N^{(k)}}$ would be an open subnormal subgroup of G such that $H' = N^{(k+1)}$ is discrete, contradicting the fact that G is assumed to be axtraordinary.

Remark 3.4 A linear p-adic Lie group G cannot be extraordinary because $L(R(G))$ would be the radical of $L(G)$ in this case (by Lemma 1.3) and thus $L(R(G)) = L(G)$ (as $L(G)$ is abelian). Since $R(G)$ is discrete (see Remark 3.3), $L(G) = \{0\}$ would follow and thus G would be discrete (contradicting the hypothesis that G is extraordinary).

Remark 3.5 For G an elementary p-adic Lie group, there is no clear relation between the dimension of G and its constructible rank. E.g., $0 = \mathrm{crk}(G) < \dim(G)$ for each non-discrete compact p-adic Lie group G. Conversely, consider $\mathbb{Q}_p \rtimes \mathbb{Z}$ with $1 \in \mathbb{Z}$ acting by the automorphism $\mathbb{Q}_p \to \mathbb{Q}_p$, $z \mapsto pz$. Then $\mathrm{crk}(G) \leq 2$ by construction and in fact $\mathrm{crk}(G) = 2 > 1 = \dim(G)$. In fact, $\mathrm{crk}(G) \neq 0$ since G is neither compact nor discrete. If we had $\mathrm{crk}(G) = 1$, then either G would be an extension $N \to G \to Q$ where N and Q are compact or discrete (each of these four possibilities can be ruled out), or G would be an ascending union of open subgroups of constructible rank 0, which therefore have to be compact (being non-discrete): Also this is impossible (since $\mathbb{Z}$ is a quotient of G).

4 Nearly simple Lie groups over other fields

Some of the above results are independent of the ground field of p-adic numbers. In this section, we let $(\mathbb{K}, |.|)$ be a complete valued field on which $|.|$ defines a non-discrete topology. We consider finite-dimensional analytic Lie groups over $(\mathbb{K}, |.|)$, as in [8]. A subgroup $S \subseteq G$ is called a Lie subgroup if it is an embedded submanifold. Such subgroups are closed, in particular.

Definition 4.1 Let G be an analytic Lie group over $\mathbb{K}$. We say that a Lie subgroup $S \subseteq G$ is *Lie subnormal* if there exists a series

$$G = G_0 \rhd G_1 \rhd \cdots \rhd G_k = S$$

of Lie subgroups G_j of G_{j-1} for $j \in \{1, \ldots, k\}$. We say that G is *nearly Lie simple* if every Lie subnormal subgroup $S \subseteq G$ is discrete or open.

The next proposition can be proved exactly as Proposition 2.2, except that closed subgroups need to be replaced with Lie subgroups.

Proposition 4.2 *Every analytic Lie group G over $\mathbb{K}$ admits a series*

$$G = G_0 \rhd G_1 \rhd \cdots \rhd G_n = \{1\}$$

of Lie subgroups G_j with G_{j-1}/G_j nearly Lie simple for all $j \in \{1, \ldots, n\}$. $\square$

Acknowledgements. I am grateful to P.-E. Caprace, C.R.E. Raja and Ph. Wesolek for discussions. The core of the manuscript was written during a visit to the Fields Institute (Toronto), partially supported by the latter.

References

[1] Bass, H., *Groups of integral representation type*, Pacific J. Math. **86** (1980), no. 1, 15–51.

[2] Baumgartner, U. and G. A. Willis, *Contraction groups and scales of automorphisms of totally disconnected locally compact groups*, Israel J. Math. **142** (2004), 221–248.

[3] Bourbaki, N., "Lie Groups and Lie Algebras, Chapters 1–3," Springer-Verlag, Berlin, 1989.

[4] Caprace, P.-E., C. D. Reid and G. A. Willis, *Limits of contraction groups and the Tits core*, J. Lie Theory **24** (2014), no. 4, 957–967.

[5] Cluckers, R., Y. Cornulier, N. Louvet, R. Tessera, and A. Valette, *The Howe-Moore property for real and p-adic Lie groups*, Math. Scand. **109** (2011), no. 2, 201–224.

[6] Glöckner, H., *Scale functions on p-adic Lie groups*, Manuscripta Math. **97** (1998), no. 2, 205–215.

[7] Margulis, G. A. "Discrete Subgroups of Semisimple Lie Groups," Springer, Berlin, 1991.

[8] Serre, J.-P., "Lie Algebras and Lie Groups," Springer, Berlin, 1992.

[9] Wang, J. S. P., *The Mautner phenomenon for p-adic Lie groups*, Math. Z. **185** (1984), 403–412.

[10] Wesolek, Ph., *Radical subgroups of totally disconnected locally compact groups*, `arXiv:1304.7522v2`

[11] Wesolek, Ph., *Elementary totally disconnected locally compact groups*, manuscript in preparation.

[12] Wesolek, Ph., *Totally disconnected locally compact groups locally of finite rank*, manuscript in preparation.

[13] Willis, G. A., *The structure of totally disconnected, locally compact groups*, Math. Ann. **300** (1994), 341–363.

Helge Glöckner, Universität Paderborn, Institut für Mathematik,
Warburger Str. 100, 33098 Paderborn, Germany

e-mail: `glockner@math.upb.de`